

## 4.1 Solving Polynomial Equations

<p><b>A Polynomial Equation</b></p> <p>A <i>polynomial equation</i> is defined as:</p> $P(x) = 0$ <p>where <math>P(x)</math> is a polynomial function.</p> <p>Note. The numbers <math>x</math> satisfying the polynomial equation are called the <i>roots</i> or the <i>solutions</i> of the polynomial equation.</p> <p>Note. The <i>roots (solutions)</i> of the polynomial equation <math>P(x) = 0</math> are the same as the <i>zeros</i> of the polynomial function <math>y = P(x)</math>.</p>	<p>Ex 1. Show that <math>x = \sqrt{3}</math> is a solution of the polynomial equation</p> $x^4 + 9 = 6x^2$
<p><b>B Grouping</b></p> <p>Some polynomial equations may be solved by grouping terms adequately.</p> <p>Note. Is not easy to see how to group terms in order to solve the equation.</p>	<p>Ex 2. Solve for <math>x</math> by grouping.</p> $8x^3 - 12x^2 - 2x + 3 = 0$
<p><b>C Integral Zeros Theorem</b></p> <p>If <math>x = b</math> is an <i>integral zero</i> of the polynomial <math>P(x)</math> with integral coefficients, then <math>b</math> is a <i>factor (divisor)</i> of the <i>constant term</i> <math>a_0</math> of the polynomial.</p> <p>Note. A <i>real zeros</i> of the polynomial function <math>P(x)</math> is also called <i>x-intercept</i> because the graph <i>touches</i> or <i>crosses</i> the <i>x-axis</i> at this number.</p>	<p>Ex 3. Solve for <math>x</math> by looking first at integral roots (solutions).</p> $(x^2 - 13)x = 15 - 3x^2$

<p><b>D Rational Zero Theorem</b></p> <p>If <math>x = \frac{b}{a}</math> is an <i>rational zero</i> of the polynomial <math>P(x)</math> with integral coefficients, then <math>b</math> is a <i>factor (divisor)</i> of the <i>constant term</i> <math>a_0</math> and <math>a</math> is a <i>factor (divisor)</i> of the <i>leading coefficient</i> <math>a_n</math>.</p>	<p>Ex 4. Solve for <math>x</math> by looking first at rational roots (solutions).</p> $3x^4 + \frac{7}{2}x^3 - \frac{2}{3}x^2 - \frac{3}{2}x - \frac{1}{3} = 0$
<p>Ex 5. Is <math>-2/3</math> a possible rational zero of <math>P(x) = 2x^3 + \dots + -8</math>? Explain.</p>	
<p><b>E Non Rational Zeros</b></p> <p>If <math>x = a + b\sqrt{c}</math> (<math>a, b, c</math> are rational numbers, <math>c &gt; 0</math>) is a zero of a polynomial with integral coefficients <math>P(x)</math>, then <math>x = a - b\sqrt{c}</math> is also a zero of this polynomial.</p>	<p>Ex 6. Solve for <math>x</math></p> $x^4 - 2x^3 - 4x^2 + 6x + 3 = 0$ <p>given that <math>x = 1 + \sqrt{2}</math> is one of its roots.</p>
<p><b>F Technology</b></p> <p>In some cases the real roots of a polynomial equation may be found only using <i>technology</i>.</p>	<p>Ex 7. How many real roots does the equation given below have?</p> $x^3 - x^2 - 1 = 0$ <p>Use technology to find (real) <math>x</math>.</p>

**Reading:** Nelson Textbook, Pages 196-204

**Homework:** Nelson Textbook, Page 204: #6ad, 7bf, 8a, 9b, 10, 13, 15, 16, 18, 19